

Evolution of the electron cyclotron drift instability in two-dimensions

Salomon Janhunen, Andrei Smolyakov, Dmytro Sydorenko, Marilyn Jimenez, Igor Kaganovich, and Yevgeny Raïtses

Citation: *Physics of Plasmas* **25**, 082308 (2018); doi: 10.1063/1.5033896

View online: <https://doi.org/10.1063/1.5033896>

View Table of Contents: <http://aip.scitation.org/toc/php/25/8>

Published by the [American Institute of Physics](#)

PHYSICS TODAY

WHITEPAPERS

MANAGER'S GUIDE

Accelerate R&D with
Multiphysics Simulation

READ NOW

PRESENTED BY

 **COMSOL**

Evolution of the electron cyclotron drift instability in two-dimensions

Salomon Janhunen,^{1(a)} Andrei Smolyakov,¹ Dmytro Sydorenko,² Marilyn Jimenez,¹ Igor Kaganovich,³ and Yevgeny Raitses³

¹University of Saskatchewan, 116 Science Place, Saskatoon, Saskatchewan S7N 5E2, Canada

²University of Alberta, 3-235 Centennial Centre for Interdisciplinary Science, Edmonton, Alberta T6G2E9, Canada

³Princeton University/Princeton Plasma Physics Lab, 100 Stellarator Rd., Princeton, New Jersey 08543-0451, USA

(Received 6 April 2018; accepted 20 July 2018; published online 14 August 2018)

The Electron Cyclotron Drift Instability driven by the electron $E \times B$ drift in partially magnetized plasmas is investigated with highly resolved particle-in-cell simulations. The emphasis is on two-dimensional effects involving the parallel dynamics along the magnetic field in a finite length plasma with dielectric walls. It is found that the instability develops as a sequence of growing cyclotron harmonics demonstrating wave breaking and complex nonlinear interactions, being particularly pronounced in ion density fluctuations at short wavelengths. At the same time, nonlinear evolution of fluctuations of the ion and electron density, as well as the anomalous electron current, shows cascade toward long wavelengths. Tendency to generate long wavelength components is most clearly observed in the spectra of the electron density and the anomalous current fluctuations. An intense but slowly growing mode with a distinct eigen-mode structure along the magnetic field develops at a later nonlinear stage enhancing the tendency toward long wavelength condensation. The latter mode having a finite wavelength along the magnetic field is identified as the Modified Two-Stream Instability (MTSI). It is shown that the MTSI mode results in strong parallel heating of electrons. *Published by AIP Publishing.* <https://doi.org/10.1063/1.5033896>

I. INTRODUCTION

Partially magnetized plasmas immersed in crossed $E \times B$ fields are used in various devices such as Hall thrusters for electric propulsion. Such plasmas are subject to a number of instabilities that affect device operation—and in particular—the level of anomalous transport that is typically found to be orders of magnitude larger than the classical (collisional) transport. The nature of the anomalous transport (mobility) is still poorly understood and has been attributed to several candidate instabilities that may interact with each other to bring about the observed levels of anomalous transport. The electron cyclotron drift instability (ECDI) driven by the electron $E \times B$ drift, and independent of any plasma gradients and collisions, has been recently actively discussed as a possible candidate.^{1–3}

In earlier works,^{4–7} electron cyclotron instabilities have been studied in relation to turbulent plasma heating by the electric current perpendicular due to the relative electron-ion drift. Electron cyclotron instability driven by ion beams was also identified as a possible source of anomalous resistivity explaining the width of collisionless shock waves, in particular in space conditions; for more recent work and references see Refs. 8–10.

In the context of the anomalous transport in Hall thrusters, the cyclotron instability driven by the electron $E \times B$ drift was studied in 1D simulations,^{3,11–13} 2D axial-azimuthal simulations,^{14–16} and 2D radial-azimuthal simulations.^{17–19} Many

of these works focused on the possibility that ECDI simply becomes the ion sound instability analogous to the case of unmagnetized plasma.^{13,20} We have shown in our previous nonlinear simulations that the transition to ion sound (which for the 1D case is only possible due to nonlinear diffusion in the short wavelength regime) does not occur²¹ and the instability is driven by the dominant $m = 1$ cyclotron resonance. We also found in our previous 1D simulations that when a larger azimuthal length is used, anomalous transport cascades to low- k modes. Curiously in Ref. 13 the authors see the emergence of a large-scale structure in their simulation when they use a larger simulation box, but reject it as an artifact.

In this paper, using highly resolved particle-in-cell simulations, we study instabilities and transport in the 2D (azimuthal-radial) geometry. Periodic boundary conditions are used in the azimuthal direction, along the $E \times B$ drift. The magnetic field is in the radial direction bound by the dielectric wall boundaries. Curvature effects of the channel are not included in this work.

In 2D geometry, a new class of unstable mode appears for finite values of the wave number k_z along the magnetic field, namely the Modified Two-Stream Instability (MTSI).²² For larger values of k_z , the unstable mode looks similar to the unmagnetized ion sound.^{5,20,23} In this paper, we study the linear and nonlinear evolution of the interacting ECDI and MTSI modes, their saturation, and associated turbulent transport for typical conditions of a Hall thruster. Our simulations demonstrate that like in the 1D case, the instability is driven by nonlinear cyclotron resonance modes that dominate the anomalous transport and that the cascade to long wavelengths observed in 1D simulations is further enhanced

Note: Invited as part of the Special Topic on Modern Issues and Applications of $E \times B$ Plasmas (Guest Editors: Jean-Pierre Boeuf and Andrei Smolyakov).

^{a)}Electronic mail: salomon.janhunen@usask.ca.

by the linear long wavelength instabilities that occur when finite k_z is allowed. Moderate values of the anomalous electron current of the order of $\Omega = \Omega_{ce}/\nu_{eff} \simeq 200$ are obtained in a nonlinear stage similar to the 1D case.²¹ An important new result is strong parallel electron heating due to the modes with a finite k_z .

This paper is structured as follows: we discuss the 2D linear regime and the unstable normal modes therein and show that they appear as expected in fully non-linear simulations in the very early part of the simulation. The early non-linear saturation processes are discussed, such as mode competition between the ECDI harmonics and apparent coupling to the modified two-stream instability (MTSI). In the development of the strong turbulence regime, we show how the MTSI compresses the ECDI wave packet and produces large fluxes to the sheath with accompanied rapid heating of the parallel temperature. We discuss the spectral cascade of the anomalous current, the features of the anomalous current as a function of the radial (parallel to magnetic field) direction, and the time evolution of the overall anomalous current. Finally, the sheath losses and decay of the plasma column in the absence of sources are discussed, and a summary follows. We discuss technical details such as numerical parameters and analysis methods in the [Appendix](#).

II. LINEAR FEATURES OF THE ELECTRON CYCLOTRON DRIFT-INSTABILITY

The electron drift cyclotron instability (ECDI) occurs in partially magnetized $E \times B$ plasma due to the significant $E \times B$ flow of electrons with respect to ions. It is convenient to discuss the characteristics of the ECDI with reference to the linear dispersion relation. We consider the electrostatic waves with $\mathbf{v}_0 = \mathbf{E} \times \mathbf{B}/B^2$ streaming of electrons across a uniform magnetic field \mathbf{B} , with unmagnetized ions, in homogeneous unbounded plasma. The two-dimensional linear dispersion equation has the form²³

$$\epsilon(\omega, \mathbf{k}) = 1 + \epsilon_i(\omega, \mathbf{k}) + \epsilon_e(\omega, \mathbf{k}) = 0, \quad (1)$$

where ϵ_e and ϵ_i are the electron and ion susceptibilities

$$\epsilon_i = -\frac{1}{2k^2\lambda_{Di}^2} Z\left(\frac{\omega}{\sqrt{2}kv_i}\right), \quad (2)$$

$$\epsilon_e = \frac{1}{k^2\lambda_{De}^2} \left[1 + \frac{\omega - \mathbf{k} \cdot \mathbf{v}_0}{\sqrt{2}k_z v_e} \sum_{m=-\infty}^{\infty} e^{-b} I_m(b) \times Z\left(\frac{\omega - \mathbf{k} \cdot \mathbf{v}_0 + m\Omega_{ce}}{\sqrt{2}k_z v_e}\right) \right], \quad (3)$$

where $b = k_y^2 \rho_e^2$, $\rho_e^2 = v_e^2/\Omega_{ce}^2$, $v_{e,i}^2 = T_{e,i}/m_{e,i}$, $\lambda_{De,i}^2 = \epsilon_0 T_{e,i}/n_0 q_{e,i}^2$, $Z(\xi)$ is the plasma dispersion function, $I_m(x)$ is the modified Bessel function of the 1st kind, $\mathbf{B} = B_0 \hat{z}$ is the magnetic field in the z -direction, and $\mathbf{E} = E_0 \hat{x}$ is the external electric field in the x -direction, so that $\mathbf{v} = \mathbf{E} \times \mathbf{B}/B^2 = v_0 \hat{y}$ is in the y direction (or, against if $v_0 = -E_0/B_0 < 0$). Then, k_z and k_y are the components of the wave vector \mathbf{k} along the magnetic field and in the $\mathbf{E} \times \mathbf{B}/B$ directions, $k \equiv |\mathbf{k}| = \sqrt{k_z^2 + k_y^2}$. Given the physical parameters, it remains to

assign k_z and k_y values; solving the dispersion relation provides us with the frequency ω , which generally gets a complex value.

For the geometry of the Hall thruster with a radial magnetic field, we define a few auxiliary quantities and relations that aid in the following discussions: $k_0 = \Omega_{ce}/v_0$, $k_z = 2\pi n_z/L_r$, $k_y = 2\pi n_y/l_\theta$, where L_r is the extent of the system in the radial (along the magnetic field) direction, and l_θ is the extent of the system in the azimuthal direction, so the quantum numbers n_z and n_y characterize the radial and azimuthal wave vectors that satisfy the boundary conditions. The local Cartesian coordinates y, z correspond to the θ, r coordinates of the coaxial Hall thruster. It is important to emphasize however that while the y -direction is periodic in our simulations, periodicity is not imposed in the radial direction. The eigenmode structure in the radial direction is formed self-consistently by the mode parallel dynamics and by the sheath effects at $z=0$ and $z=L_r$. The ensuing mode structure will be discussed below. In this work, we assume the gap to be a straight box, for simplicity.

In the limit of cold ions where $\omega > kv_i$, the ion response becomes

$$\epsilon_i = -\frac{\omega_{pi}^2}{\omega^2}, \quad (4)$$

where $\omega_{pi}^2 = e^2 n_0/\epsilon_0 m_i$ is the ion plasma frequency.

In the one-dimensional case, $k_z \rightarrow 0$, $k = k_y$, the dispersion equation (1) takes the form

$$\epsilon_e = \frac{1}{k^2 \lambda_{De}^2} \left[1 - \exp(-k^2 \rho_e^2) I_0(k^2 \rho_e^2) - 2(\omega - kv_0)^2 \sum_{m=1}^{\infty} \frac{\exp(-k^2 \rho_e^2) I_m(k^2 \rho_e^2)}{(\omega - kv_0)^2 - m^2 \Omega_{ce}^2} \right]. \quad (5)$$

The form of Eq. (5) emphasizes the role and the interaction of different cyclotron harmonics. Note that there is no resonance for the $m=0$ harmonic, while all higher harmonics with $m=1, 2, \dots$ are resonant at $(\omega - kv_0)^2 = m^2 \Omega_{ce}^2$. In the cold electron limit $T_e \rightarrow 0$, only $m=0$ and $m=1$ harmonics contribute and the dispersion relation reduces to the Buneman magnetized plasma instability driven by transverse current²⁴

$$1 - \frac{\omega_{pi}^2}{\omega^2} - \frac{\omega_{pe}^2}{(\omega - kv_0)^2 - \Omega_{ce}^2} = 0. \quad (6)$$

The instability is a result of reactive coupling of the electron (Doppler shifted) upper hybrid mode $(\omega - kv_0)^2 = \omega_{pe}^2 + \Omega_{ce}^2$ with the short wavelength ion oscillations $\omega^2 = \omega_{pi}^2$. In the long wavelength low frequency limit, $(\omega, kv_0) < \Omega_{ce}$, Eq. (6) describes the lower-hybrid modes, $\omega_{LH}^2 = \Omega_{ci} \Omega_{ce}$. The contribution of higher $m > 1$ harmonics (which are absent for $T_e = 0$) grows with electron temperature and has the maximum at shorter wavelengths $k^2 \rho_e^2 \simeq 1$ due to the $\exp(-k^2 \rho_e^2) I_m(k^2 \rho_e^2)$ factors. The temperature effects also add dispersion to the lower hybrid modes $\omega^2 = \Omega_{LH}^2 (1 + k^2 \rho_e^2)$, which eventually becomes the high frequency ion sound for $k^2 \rho_e^2 \gg 1$, ω^2

$= \Omega_{LH}^2 k^2 \rho_e^2 \simeq k^2 c_s^2$. In this limit, the contribution of the cyclotron harmonics decreases with $k\rho_e$: $\exp(-k^2 \rho_e^2) \times I_m(k^2 \rho_e^2) \rightarrow 1/(k\rho_e)$, so that the real part of the electron susceptibility becomes

$$\epsilon_e \simeq 1 / \left(k^2 \lambda_{De}^2 \right), \quad (7)$$

and Eq. (1) produces the ion sound mode

$$\omega^2 = k^2 c_s^2 / \left(1 + k^2 \lambda_{De}^2 \right). \quad (8)$$

The imaginary part in ϵ_e (neglected so far) originates in the series of cyclotron resonances. In the limit $k^2 \rho_e^2 \gg 1$, the infinite series of cyclotron resonances can be summed resulting in the imaginary contribution equivalent to the pole contribution $1/(\omega - kv_0)$ as for the case of unmagnetized electrons. Thus, even for strictly perpendicular propagation, in the $k^2 \rho_e^2 \gg 1$ limit, one has the ion sound instability as in the unmagnetized plasma case. This case is directly related to the resolution of the Landau-Bernstein paradox: the sequence of Bernstein modes which are undamped in magnetized plasma results in collisionless Landau damping when $B \rightarrow 0$.

Another kind of instability occurs near the resonances $(\omega - kv_0)^2 \simeq m^2 \Omega_{ce}^2$. This is a strong (fluid) reactive instability due to coupling of the ion and electron modes,²⁵ facilitated by the Doppler shift. For cold electrons, only the $m = 1$ exists, resulting in the Buneman instability described by Eq. (6). For finite T_e , all higher modes with $m > 1$ are present. In our previous work, it was shown that in the 1D case a set of modes with higher m are excited, but eventually, via the linear (due to electron heating) and nonlinear effects, a dominant $m = 1$ strongly coherent cnoidal wave appears.²¹ The cyclotron resonance nature of the mode, defined by the condition $\omega < k_y v_0 \simeq \Omega_{ce}$, extends far into the nonlinear stage. Similar results were also obtained in other simulations relevant to space plasma conditions.⁸

In 2D where plasma motion along the magnetic field is present, new regimes become possible due to finite values of the k_z . There have been a number of studies of the full linear dispersion equation with (2) and (3), for example see Refs. 5, 20, 23, and 26. One of the results of these studies is that for sufficiently large values of the parallel wave vector k_z , the solution of the dispersion relation (1) produces a mode which is close to the ion sound instability in unmagnetized plasma driven by the electron beam with v_0 velocity.

The general behavior of the growth rate of the instability is shown for several values of the k_z parameter in Fig. 1 as a function of the azimuthal wave vector k_y . We solve the dispersion relation numerically in Python²⁷ using the technique described in Ref. 20, where the solution is obtained through a fixed point iteration using the relative error as a stopping condition. We use the convergence condition that $|1 - \omega_{i+1}/\omega_i| < 10^{-6}$ for $i \geq 15$. The SciPy²⁸ Faddeeva function is used to get good numerical accuracy of the plasma dispersion function for a wide range of arguments. We show the first four roots obtained in the (k_z, k_y) phase plane in Fig. 2 for a 10eV Xenon plasma with $n_e = 10^{17} \text{ m}^{-3}$, $B_0 = 0.02 \text{ T}$, and $E_0 = 20 \text{ kV/m}$, which are typical Hall thruster parameters.




FIG. 1. Growth rate of instabilities for $k_z \lambda_{De}$ of 0.01, 0.02, 0.04, and 0.08, respectively, found from the full dispersion equation (1)–(3). The first root from left is the MTSI ($m = 0$), and subsequent roots are from $m = 1, 2, 3, \dots$ ECDI resonances. Parameters for this figure are: $v_0 = -369 c_s$, $v_{Te} = 489 c_s$, $\Omega_{ce} = 96.5 \omega_{pi}$, and $k_0 \lambda_{De} = 0.262$.

The normalization scheme for the dispersion relation solver code is the same as defined in Ref. 20, where frequencies are normalized to ω_{pi} , velocities to c_s , and lengths to λ_{De} . The physical parameters given above are also used as the initial state of nonlinear simulations presented in this paper, unless otherwise noted. The Python code is also used for solving the dispersion relation in a simple case of $T_e \rightarrow 0$, as shown in Fig. 3 where the limit of the full dispersion relation is presented for $k_z \lambda_{De} = 0.005$.

For small values of the k_z , one can see in Fig. 1 the fluid type reactive instability peaks near the resonant values of $k_y = m \cdot k_0$. For larger values of k_z , the fluid resonance is broadened by the thermal effects and for $k_z v_{Te} \geq \omega$, the resonance becomes kinetic, resulting in the smooth curve corresponding to the kinetic resonance instability of the ion sound in unmagnetized plasma.²³

The regimes for different values of k_z can be seen from the dispersion equation (1). In the limit $k_z \rightarrow 0$, $\xi_m \gg 1$, and using $Z(\xi_m) \rightarrow -1/\xi_m$, the terms involving plasma dispersion functions result in cyclotron resonances




FIG. 2. Phase space plot of growth rate $\gamma(k_z, k_y)/\omega_{pi}$ for the full dispersion relation. Notice the modified two-stream instability (MTSI) in the sub-cyclotron low- k region, indicating possible instability for simulations which are able to accommodate the mode in the azimuthal direction. Parameters for this figure are: $v_0 = -369 c_s$, $v_{Te} = 489 c_s$, $\Omega_{ce} = 96.5 \omega_{pi}$, and $k_0 \lambda_{De} = 0.262$.




FIG. 3. Comparison of the growth rate from a 2D simulation and a 1D simulation with 40 000 particles/cell, using $T_e = 0.001$, with the solution of the linear dispersion in the $T_e \rightarrow 0$ limit. Parameters for this figure are: $v_0 = -36.9 \cdot 10^3 c_s$, $v_{Te} = 489 c_s$, $\Omega_{ce} = 96.5 \omega_{pi}$.

$$\frac{\omega - \mathbf{k} \cdot \mathbf{v}_0}{\sqrt{2}k_z v_e} Z(\xi_m) \rightarrow -\frac{\omega - \mathbf{k} \cdot \mathbf{v}_0}{\omega - \mathbf{k} \cdot \mathbf{v}_0 + m\Omega_{ce}}, \quad (9)$$

$$\xi_m = \frac{\omega - \mathbf{k} \cdot \mathbf{v}_0 + m\Omega_{ce}}{\sqrt{2}k_z v_e}. \quad (10)$$

This is the regime of the fluid (reactive) instability due to coupling of the ion sound and ion modes²⁵ which occurs for cold plasma (and in the limit of $k_z \rightarrow 0$).

The thermal broadening of the resonance and the transition to the kinetic ion sound instability can be clearly seen in the example when only $m = 0$ is retained in the sum (3)

$$1 - \frac{\omega_{pi}^2}{\omega^2} + \frac{1}{k^2 \lambda_{De}^2} \left[1 + \frac{\omega - \mathbf{k} \cdot \mathbf{v}_0}{\sqrt{2}k_z v_e} e^{-b} I_0(b) Z\left(\frac{\omega - \mathbf{k} \cdot \mathbf{v}_0}{\sqrt{2}k_z v_e}\right) \right] = 0. \quad (11)$$

The first three terms in this expression describe the ion sound mode (8). Considering the last term as a small perturbation in Eq. (11), one gets the so called modified ion sound instability.^{26,29} Similar resonance broadening occurs for higher m resonances, so that the resonant fluid type instability ($k_z \rightarrow 0$) is replaced by the kinetically driven mode for a finite k_z . This behavior is illustrated in Figs. 1 and 4, where the mode frequencies and growth rates are calculated retaining only individual terms with different m of the Bessel function series (lower panel) and partial sum up to the m th order (upper panel). Note that the real part of the mode frequency for individual m and partial sum m modes always remains close to the ion sound mode frequency from Eq. (8), see Fig. 4. For sufficiently large k_z , the summation of several components in the Bessel series illustrates the transition to unmagnetized ion-sound instability as shown in Fig. 4, where the solutions of the partial sum slowly converge to the unmagnetized curve as terms are added.

III. MODIFIED TWO-STREAM INSTABILITY AND MODIFIED TWO-STREAM BUNEMAN INSTABILITY

The discussion in Sec. II remains silent on one important feature of the instability with finite k_z : namely on the regime

of the Modified-Two-Stream Instability (MTSI)^{22,30,31} where the effect of the electron parallel motion is typically considered in the context of the following dispersion equation:³⁰

$$1 - \frac{\omega_{pi}^2}{\omega^2} - \frac{\omega_{pe}^2 k_z^2}{(\omega - k_y v_0)^2 k^2} + \frac{\omega_{pe}^2 k_y^2}{\Omega_{ce}^2 k^2} = 0. \quad (12)$$

It is instructive to analyze the nature of this equation starting from the electron response in the form of Eq. (7) which together with Eq. (4) results in the ion sound mode. The electron susceptibility in the form of Eq. (7) is equivalent to the Boltzmann response for the perturbed electron density

$$\tilde{n}_e = \frac{e\tilde{\phi}}{T_e} n_0. \quad (13)$$

This expression follows from the parallel electron balance in neglect of the electron inertia:

$$0 = en\nabla_{\parallel}\tilde{\phi} - T_e\nabla_{\parallel}\tilde{n}. \quad (14)$$

The electron inertia however can be neglected in the parallel momentum balance only when the condition $\omega < k_z v_e$ is satisfied. In the presence of the strong transverse electron flow v_0 , apparent mode frequency has to be modified due to the Doppler shift: $\omega \rightarrow \omega - k_y v_0$. When the condition $\omega - k_y v_0 < k_z v_e$ is violated, the electron inertia terms have to be included. In this case, the electron continuity and momentum balance equations

$$-i(\omega - k_y v_0)\tilde{n} + ik_z n_0 v_{\parallel} = 0, \quad (15)$$

$$-im_e(\omega - k_y v_0)v_{\parallel} = ik_z \tilde{\phi}, \quad (16)$$

result in the electron density response in the form

$$n_e = -\frac{ek_z^2 \tilde{\phi}}{m_e(\omega - k_y v_0)^2}. \quad (17)$$

Note that when $(\omega - k_y v_0) \simeq k_z v_e$, the assumption of the isothermal electrons becomes invalid, and the only consistent approximation in the fluid theory is to assume $T_e = 0$, so the pressure term in (16) was omitted. In this equation, the third term corresponds to the electron density perturbation from Eq. (17) and the last term is due to the electron inertial perpendicular current. It is worth noting that Eq. (12) is not fully consistent. The second term in this equation is obtained under ordering $(\omega - k_y v_0) \simeq k_z v_e$, while the last term is obtained with the low frequency approximation $(\omega - k_y v_0) \ll \Omega_{ce}$. The latter may not be satisfied for some applications such as Hall thrusters. A more accurate dispersion equation is obtained from Eqs. (1) and (3) by taking a rigorous limit $T_e \rightarrow 0$ which yields the relation

$$1 - \frac{\omega_{pi}^2}{\omega^2} - \frac{\omega_{pe}^2 k_z^2}{(\omega - k_y v_0)^2 k^2} - \frac{\omega_{pe}^2 k_y^2}{((\omega - k_y v_0)^2 - \Omega_{ce}^2) k^2} = 0, \quad (18)$$

which includes both the modified two-stream instability and the upper hybrid Buneman instability. In the following, we




FIG. 4. Growth rate obtained from partial sums of the full dispersion relation (1); top row: terms up to $I_{\pm k}$ included solving for ω and γ ; bottom row: only terms $I_{\pm k}$ of the sum included while solving for ω and γ . The solution from the unmagnetized dispersion relation is included (labeled as IS); partial sum including terms up to $m = 50$ is virtually indistinguishable from the unmagnetized solution. Parameters for this figure are: $v_0 = -369 c_s$, $v_{Te} = 489 c_s$, $\Omega_{ce} = 96.5 \omega_{pi}$.

will call this case the Modified Buneman Two-Stream Instability (MBTSI).

The MBTSI regime is of particular importance as a finite value of k_z results in the long wavelength instability at small $k_y/k_0 \ll 1$, well below the cyclotron resonances with $k_y \simeq m \cdot k_0$. This long wavelength mode, the leftmost peak in Fig. 1, also shown in Fig. 5, has a small growth rate, but as discussed in Sec. V, turns out to be important in the nonlinear saturation regime enhancing the tendency toward long wavelength condensation.

In our 2D simulations, which are bounded in the z -direction, it is observed generally that up to the non-linear regime the instability grows uniformly everywhere, with $k_z \simeq 0$, and only after the non-linear regime is reached does the wave like structure develop in the parallel direction. This can be explained by the fact that the linear growth rates increase monotonically towards $k_z \rightarrow 0$, while the sheath allows fluctuations to extend up to the boundary by “insulating” the perturbations from the wall.

Generally in the literature, the ECDI instability was considered²⁰ for typical values of the parallel wave length of $k_z \lambda_{De} \simeq 0.01$ – 0.09 corresponding to a parallel wave length of $\lambda = 7.4$ – 0.8 cm, for typical Hall-effect thruster parameters. It




FIG. 5. Close-up of the modified two-stream instability (MTSI) growth rate γ/ω_{pi} given as a function of $\sin(\alpha) = k_z/k$ and k/k_0 . Parameters for this figure are: $v_0 = -369 c_s$, $v_{Te} = 489 c_s$, $\Omega_{ce} = 96.5 \omega_{pi}$, $k_0 \lambda_{De} = 0.262$.




FIG. 6. Growth rate and frequency for ECDI instabilities for $k_z \lambda_{De} = 0.005$ at $T_e = 10$ eV. Growth rates as obtained from the 2D simulation (Fig. 8) are indicated by the symbols. Note the existence of the MTSI below the first cyclotron resonance. Refer to Table I for the values. Parameters for this figure are: $v_0 = -369 c_s$, $v_{Te} = 489 c_s$, $\Omega_{ce} = 96.5 \omega_{pi}$, $k_0 \lambda_{De} = 0.262$.

could be misleading, however, to estimate k_z and relevant dynamic regimes of the ECDI based on full wave lengths in a Hall-effect thruster gap. It has been noted earlier^{32–35} that in a bounded plasma the sheath effects allow the wavelength along the magnetic field to be much longer than would otherwise follow from the geometrical constraint $k_z = 2\pi/L_r$, where L_r is the plasma width between the boundaries. In our 2D simulations, we observe that the parallel structure along

the magnetic field is consistent with a half-wavelength fit between the boundaries. This corresponds to $k_z \lambda_{De} = 0.00434$. We used this value for calculations of the linear growth rates in Figs. 3 and 6.

IV. GROWTH AND SATURATION OF THE CYCLOTRON HARMONICS OF ECDI AND MTSI MODES

PIC2D is a 2D3V PIC code developed by Dmytro Sydorenko at the University of Alberta based on his earlier 1D3V PIC code, EDIPIC.³⁶ The initial state of the PIC2D simulation is a v_0 -shifted Maxwellian distribution for the electrons and a stationary Maxwellian for the ions. The initial state is quasi-neutral and homogeneous, equipped with the typical Hall-effect thruster parameters presented in Sec. II (and Appendix A). The magnetic field lines are terminated at a dielectric boundary, so a sheath potential develops soon after time starts running. The simulation box is $L_r = 53.8$ mm by $l_\theta = 13.45$ mm, and the numerical parameters are detailed in Table II in Appendix A. Even though the code allows to do so, no collisions were used in our simulations.

Chronologically, the evolution of the simulations goes as follows (see Figs. 7 and 8). First, in the linear-like stage the fastest growing modes—the ECDI with $m = 3–5$ —grow and introduce some heating to the electrons primarily in the perpendicular direction. Due to heating and nonlinearities, the lowest $m = 1$ resonance becomes the ECDI mode with the largest energy content, even though linearly it is the slowest growing mode. The resonance condition $k_y = m\Omega_{ce}/v_0$ gives




FIG. 7. Time slices of the ion density fluctuation $n_i - \langle n_i \rangle_y(z, t)$, showing growth and saturation of the predominant $m = 1$ cyclotron mode followed by the development of the structure parallel to the magnetic field.




FIG. 8. Logarithmic amplitude of the azimuthal electric field energy $|E_y|^2$, showing stages of linear growth and early nonlinear saturation of the cyclotron and MTSI modes. Linear least-squares fitting has been used to obtain values of the growth rates for each mode (labeled by their k values wrt/ k_0), given in Table I.

with our choice of $l_\theta = 13.45$ mm the quantum numbers $n_y = m \cdot 7.534$, but due to kinetic effects the maximum growth rates are found at higher values of n_y . It is observed consistently with the linear dispersion relation that the maximum growth rates for the m cyclotron resonances correspond to $n_y(m = 1, 2, 3) = \{10, 17, 24, 31, \dots\}$ as ECDI modes. For higher resonances, the up-shift in k_y is lower, diminishing the gap between maximum growth rates of m -resonances to $\Delta n_y = 7$. The mode amplitudes are shown in Fig. 7, and growth rates are given in Table I. As can be also seen from Fig. 6, these locations are very close to the maximum growth rates obtained from the linear dispersion relation. After the $m = 1$ and $m = 2$ modes saturate, the MTSI mode starts growing, suggesting non-linear feedback between the modes. During the growth of the MTSI, mode competition between $m = 1$ and $m = 2$ ECDI modes is observed. The MTSI mode grows, heating the electrons predominantly in the parallel direction due to the parallel electric field. Heating in the parallel direction results in enhanced losses to the sheath and saturation of the MTSI. The ECDI modes stay at their saturated level that was established earlier, but after the saturation of the MTSI the $m = 1$ and 2 modes grow by 5%–10%, while $m = 3$ and 4 modes lose energy correspondingly. At this stage, we observe the saturation of the anomalous axial current, as shown in Fig. 16.

Linear growth rates may be determined directly from simulation data when the runs are performed with good enough resolution. The procedure for finding the linear growth rates from the spectrogram of a simulation is outlined

TABLE I. The wave length and growth rate of unstable modes observed in the linear stage simulations as shown in Figs. 8 and 6.

Symbol	n_y	k/k_0	γ/ω_{pi}
▲	1	0.1325	0.4957
+	10	1.3248	0.7133
▼	17	2.2521	0.7629
•	24	3.1794	1.2568
■	31	4.1068	1.1628

in Appendix A. The linear and early non-linear stage of the evolution of individual mode resonances is shown in Fig. 8, where we also provide the values for the normalized wave number from the linear fits that can be made to the modes. The values obtained from simulations are given in Table I and also plotted for the comparison with the linear dispersion relation solutions in Fig. 6. As can be seen, the growth rates are of the right magnitude, although depressed due to the short time of growth available for fitting. The MTSI mode (first peak) is well represented though, showing the importance of good statistics. The MTSI peak is well represented because it grows to a higher amplitude, giving a larger range for least-squares fitting. This explains why the ECDI growth rates are generally slightly underestimated by the 2D simulations; the mode energy for the ECDI modes has only four periods of growth (before nonlinear stage), whereas the MTSI has more than ten periods for fitting (see Fig. 8). Part of the ECDI growth curve is affected by early nonlinear saturation processes, decreasing the apparent growth rate.

To emphasize this point, we ran a case for $T_e \approx 0$ with the 2D code, and a case with 1D version of the code using very good statistics (Fig. 3) that gives a remarkably good value for both the growth rate and frequency of the Modified Buneman Two-Stream instability [Eq. (18)].

A curious feature of the ECDI/MTSI energy in Fig. 8 is how the growth of the MTSI commences only after the ECDI modes have saturated, and how the energy of the ECDI modes remains relatively constant, while the MTSI is growing. This emphasizes that the modes are not in fact independent, but are nonlinearly coupled instead. It is observed that in the sheath-bounded radial direction the mode assumes a half-wave pattern with $n_z = 1/2$, whereas in the azimuthal (periodic) direction the mode is a full wave with $n_y = 1$. If we use this observation to prescribe $k_z = \pi/L_r$, we get Fig. 6 from the dispersion relation, and our MTSI peak aligns well with the peak of the MTSI in $k_y = 2\pi/l_\theta$, and the fit to ECDI modes with $n_y = \{10, 17, 24, 31, \dots\}$ is also satisfactory.

There are 3D simulations that indicate the existence of long-wavelength modes very similar to those found in our simulations,³⁷ suggesting that the presence of the MTSI is a robust feature even in full Hall-effect thruster geometry.

V. NONLINEAR SPECTRA OF ION AND ELECTRON DENSITY: SHORT WAVELENGTH FEATURES

We have earlier noted that the cyclotron resonances drive strongly coherent cnoidal type waves which are limited by saturation through ion dynamics,²¹ while the waves retain their cyclotron resonance characteristics far into the simulation even with significant electron heating and nonlinear interactions between modes.

In 2D, the dispersion relation admits a long wavelength instability that is absent in 1D, namely the modified two-stream instability of Sec. II. The latter mode is observed to modify the nonlinear dynamics compared to the 1D case, resulting in significantly faster evolution of the long wavelength components, and a lower energy content (or, fluctuation level) is retained in the ECDI modes.




FIG. 9. Ion and electron density fluctuation levels over the whole simulation volume; standard deviation and maximum.

After the early nonlinear stage, the convective nonlinearity compresses/expands the ECDI in the hills/troughs of the now-dominant MTSI mode (Fig. 10), causing jet-like injection into the sheath from the compressed maxima (particularly evident in the 4th figure of Fig. 7) with accompanying faster decay of the plasma profile. Transient large fluctuations in the electron density predominantly in the parallel direction are observed to originate from the sheath at this stage, also seen as parallel ripples in the ion density, which likely act as a relaxation mechanism. At this stage (around $1 \mu\text{s}$ into the simulation), the linear mode characteristics become a poor descriptor of the system: electron density fluctuations do not significantly increase, but ion density fluctuations grow (see Figs. 9–11), and group velocity of the wave packet increases significantly. Feedback between the ECDI- and MTSI-scale modes is apparent, with expansion and compression of the wave crests as the wave packet progresses.

An important feature of the nonlinear dynamics observed in our 2D simulation is the difference in the behavior of the ion and electron density. After heating, electron fluctuations become fairly uninteresting (and low-amplitude), but the ion




FIG. 10. Stages of the nonlinear development of the ion density fluctuations at four time slices. A large scale mode is formed, the wave crest compressing the EC waves, merging of the peaks with shift to lower k and reshaping of the wave packet to a more triangular form. Plots are from $r = 5 \text{ mm}$.

density exhibits a wealth of non-linear phenomena. This difference is especially apparent in the short wavelength $k_y \lambda_{De} < 1$ part of the spectrum. The ECDI still remains the dominant mode of energy injection into the short wavelength ion-sound fluctuations, whose frequency approaches ω_{pi} in this limit, but strongly modified by signatures of nonlinear wave breaking³⁸ due to ion dynamics. In the strong turbulence state, the ion density fluctuations in the azimuthal direction become cnoidal-like and are led by a wave with a very sharp peak of positive amplitude, after which a train of crests of decreasing amplitude follow. As apparent from Fig. 11, where a radial section of the plasma at 5 mm is shown, the crests do not propagate as much as exchange energy through elastic-like collisions (akin to soliton collisions), so the amplitude maximum travels at a higher speed than the individual crests do, as may be the case for envelope solitons.³⁹ Similar features were




FIG. 11. Left: Azimuthal ion density fluctuations as a function of time. Individual wave crests propagate at different phase velocities with respect to the wave packet, exchanging energy with one another. Right: Azimuthal electron density fluctuations as a function of time. After saturation, electron density assumes a smooth profile. The volume-averaged root-mean-square fluctuation level is roughly half of the ion fluctuation level. The figures are scaled the same way, $n_0 = 10^{17} \text{m}^{-3}$ to fix units although $\delta n = n - \langle n \rangle_y(z, t)$, and both are measured at $z = 5 \text{ mm}$.

observed in 1D simulations as well, although realized after a significantly longer time of simulation.

VI. PARALLEL ELECTRON HEATING DUE TO MTSI

A new feature with respect to our earlier 1D simulations is the rapid parallel heating in the 2D simulations, observed to occur in the same pattern as the MTSI mode. It is well-known that the MTSI is an effective heating mechanism for electrons along the magnetic field²² and the heating is the likely saturation mechanism for the MTSI, as larger parallel temperature will induce large losses of high energy electrons into the sheath.

After saturation of parallel temperature $T_{e\parallel}$, electron heating in the perpendicular direction catches up to parallel heating (which has saturated through sheath losses), after which the state of strong turbulence is reached (see Fig. 12). Even in this stage, the ECDI mode structure is prominent in density spectra and is clearly affected into a “wave street” (alternating maxima and minima) characteristic to the MTSI mode, although the large scale structure is less apparent. These features are unchanged in simulations with half time-step, ruling out the possibility of numerical heating. The heating profile is shown in Fig. 13, clearly indicating the MTSI mode as the cause of parallel heating (and to a lesser degree, perpendicular). The growth of the MTSI mode terminates when the $T_{e\parallel}$ growth terminates in Fig. 8.

VII. SPECTRAL CASCADE DOWN TO LONG WAVELENGTHS

In the previous work,²¹ linearly stable long wavelength components attributable to nonlinear processes were observed to arise in the 1D system. Energy cascade toward long wavelengths is also observed in 2D simulations, with the difference that in 2D the saturation and nonlinear stage is reached much more quickly due to the presence of the linearly unstable long wavelength MTSI mode. A salient feature observed in Fig. 8 is that the MTSI becomes active only once the cyclotron modes have saturated, and the cyclotron modes respond to the saturation of the MTSI by resuming growth. This enhances the modulational nonlinear coupling through a faster linear response. The ion density k_y spectrum




FIG. 12. Anisotropic heating of the $T_{e\perp}$ and $T_{e\parallel}$ components. Parallel heating is due to the MTSI mode. The curves show the volume averaged parallel and perpendicular temperatures.




FIG. 13. Spatial structure of the perpendicular $T_{e\perp}$ and parallel $T_{e\parallel}$ electron temperatures.

at $r = 1.35$ cm, shown in Fig. 14, clearly demonstrates the progression towards lower- k modes through inverse cascade, and emergence of the turbulent spectrum soon after the MTSI mode saturates.

Cascade to low- k is even more drastic in electron density, where the $m > 1$ resonances become all but absent. This process coincides with the generation of low- k components in the anomalous current and subsequent growth of net current. For the anomalous current, a similar spectrum may be




FIG. 14. Evolution of the azimuthal ion density (top) and electron density (bottom) k -spectra over time at $L/4$ of the simulation, plotted as $\log_{10} n_{i,e}(k_y)$. Discrete peaks occur at the $\omega - kv_0 - m\Omega_{ce}$ resonances, and the lowest peak (after $0.5 \mu s$) is the MTSI mode around $k_y = 2\pi/l_0$, or $n_y = 1$.




FIG. 15. Evolution of the anomalous current spectrum $\log_{10}|\tilde{J}_x(k_y)|^2$ over time at $z = L_r/4$ of the simulation. It is notable that the $n_y = 1$ mode dominates soon after nonlinear saturation of the modes.

obtained (Fig. 15) illustrating the initial cascade towards low- k during the saturation of the ECDI modes.

The present simulations encompass only a small fraction of the total Hall-effect thruster gap in the azimuthal direction, due to their computational expense, and even the longest wavelength modes remain short compared to the total gap circumference.

VIII. ANOMALOUS CURRENT

The axial anomalous current can be calculated from PIC2D as the relative charge flux between ions and electrons. The electric field exists only in the (y, z) -plane, but the axial anomalous current arises because of the cyclotron rotation in the (x, y) -plane. In particular, azimuthal electric field fluctuations (in y -direction) result in the electron displacement in the x -direction. The anomalous current J_x obtained as the total charge flux in the x -direction is calculated directly from particle motion in the code.

The inverse cascade is particularly apparent in the spectrum of the anomalous current (Fig. 15). Another feature of the spectrum is the dominance of the $n_y = 1$ mode that corresponds to the MTSI scale length. The mode also has the same radial envelope as the current profile (Fig. 17).

The spatial structure of the anomalous current in 2D has some special features which are not possible in 1D




FIG. 16. Evolution of the total anomalous current density J_z averaged over the simulation region. The anomalous current is normalized to the inverse of the Hall parameter.¹³ The value obtained from 1D presented in Ref. 21, 1/164.




FIG. 17. Total anomalous current components $n_y = \{0, 1\}$ for the nominal case. The total current is positive definite (blue curve), but the dominant $n_y = 1$ component has a sheath-bounded half-wave radial profile corresponding to $n_z = 1/2$.

simulations. The net axial current is associated with the $k_y = 0$ component, which is plotted in Fig. 16 as a function of time. Like in the 1D simulations, we observe that the anomalous current experiences an overshoot in the saturation stage and settles down to a much lower level. The anomalous current spectrum is dominated by low- k modes, particularly by the lowest mode available to the system in the azimuthal direction, but also has a strong radial variation that is illustrated in Fig. 17. Even though the MTSI creates a large transient in the total anomalous current, after the strong turbulent regime is established the net anomalous current falls to levels that are similar to those observed in 1D simulations. The long-wavelength features in the parallel direction do not contribute to the total volume-averaged current, but could perhaps be observed in localized measurements as large alternating axial jets in the anomalous current.

IX. SHEATH LOSSES AND DECAY OF THE PLASMA COLUMN

The plasma in these simulations is bounded by dielectric boundaries in the z -direction. Charge accumulation on the dielectric surface was allowed and the displacement current in the dielectric with $\epsilon = 4.5$ was taken into account using the model of Ref. 40.

A sheath (potential) develops very quickly in the simulation before any modes have had time to grow. After having formed, the sheath does not in fact expand much into the plasma (evident in Fig. 18), but once the MTSI saturates we observe a faster decay rate of the plasma, likely due to the rapid electron heating during wave breaking. At $0.5 \mu\text{s}$, the transition to a different regime is apparent particularly in Fig. 19, where the rate of sheath losses increases drastically and the pre-sheath begins to rapidly expand into the plasma. This is the reason why our simulation box is larger than typical Hall-effect thruster gaps in the magnetic field direction—without sources the expansion of the pre-sheath would happen too soon, and physics present in a steady-state thruster gap would be dynamically obscured.




FIG. 18. Evolution of the azimuthal mean of the electron density as a function of time.

Because the simulations presented in this paper do not have sources, it is an important question whether the decay due to sheath losses is significant enough to alter interpretation of the simulation results, and if so, in what manner. Decay of the density profile is shown in Figs. 18 and 19, where the electron density profile in the magnetic field direction and the total volume averaged electron density are shown as a function of time. Decay of the plasma column and its effect on the modes are evident from Fig. 7, where the modes terminate in the z -direction where the density drop-off of the pre-sheath begins, except at later stages for a more gradual profile. Even at $2 \mu s$, it appears that the nonlinear mode structure has remained largely intact (exhibiting characteristics of the cyclotron mode) although the plasma column has already decayed significantly (40%). The profile is largely unaffected by pre-sheath expansion up to $1 \mu s$ as seen from Fig. 18, and about 10% of the electron density is lost so far.

Hence, we contend that the plasma column decay appears to have fairly little effect on the modes. Larger effects could be expected from the increase in electron temperature, which tends to modify the linear spectrum towards lower cyclotron resonances, as well as with secondary electrons (not included) that may even reverse the sheath. It is therefore likely that the robust features observed in these




FIG. 19. Evolution of the volume-averaged electron density over time.

simulations would remain in the presence of sources and sinks too if a steady state can be achieved. There are some inherent difficulties in using sources (axial feeding and ionization) to achieve a steady-state even in 1D simulations, because of feedback between the modes and the sources (heating for ionization, transport for axial feeding).

Particularly in contrast to earlier work by Héron *et al.*,¹⁹ it is somewhat disappointing that the parallel heating mechanism has not been investigated to a greater degree in the literature. This makes it difficult to assign relative importance to the inclusion of secondary emission.

X. SUMMARY AND CONCLUSION

In earlier numerical studies of the electron cyclotron instability in 1D geometry Refs. 6 and 41, it was found that the linear (exponential growth) stage of the fast beam cyclotron instability is saturated due to nonlinear turbulent broadening, which smears out the cyclotron resonances and the instability transitions into much slower ion-sound instability much like in the ordinary unmagnetized plasma. The authors of Refs. 4 and 42 have performed analogous numerical studies for similar conditions and maintained that many properties of the observed instabilities are unlike those of the ion-sound mode of unmagnetized plasma. The apparent controversies from these simulations with regard to the importance and role of nonlinear electron diffusion and electron and ion trapping, as well as the role of the finite ion temperature, have been discussed and contrasted at some length in two complementary papers, Refs. 43 and 44.

The electron cyclotron drift instability driven by the electron current has been suggested as a possible candidate for enhanced electron transport in Hall thruster in more recent Refs. 11 and 14. The authors of Refs. 2, 3, 12, 13, 15, and 17 have performed a number of numerical simulations of ECDI and mostly concluded that the instability is analogous to the ion sound instability in the absence of the magnetic field. Our recent 1D simulations²¹ performed with higher resolution and with a longer azimuthal simulation box have not confirmed this conclusion. In our simulations, we found that the criteria for the nonlinear resonance broadening and the destruction of cyclotron resonances is not satisfied and the instability proceeds as a coherent mode driven at the main cyclotron resonance (of the reactive fluid type) $k_y v_0 = \Omega_{ce}$ well into the nonlinear stage. A strong inverse energy cascade towards the longer wavelength was identified and it was shown that the anomalous current is dominated by the long wavelength modes.

In the 1D case, the transition to the ion sound regime may only occur due to the nonlinear resonance broadening and/or collisions. The role of numerical collisions was also discussed in Ref. 44. Numerical noise as well as particle re-injection (to limit the energy growth and mimic the axial direction) may also work as collisions. As it was discussed in Sec. II above, in 2D geometry, when the direction along the magnetic field is resolved, the resonance thermal broadening due to a finite $k_z c_s$ may also facilitate the transition to the ion sound regime. However, arbitrary fluctuations are allowed at a sheath boundary, facilitating access to the $k_z \approx 0$ regime

where linear growth is largest. In this paper, we have discussed the normal modes that are obtained from different limits of the full electrostatic dispersion relation and compared them with the simulation results. Using the value of the effective k_z obtained from simulations, we have found that the linear dispersion relation predicts the discrete cyclotron resonance driven modes and the long wavelength MTSI modes which were also confirmed in the simulations. It appears therefore that for our parameters the full ion sound regime appears not to be realized, but instead a more effective regime of discrete cyclotron resonance instabilities and the long wavelength MTSI occur. The magnetic field remains to be a defining feature of these regimes (contrary to the unmagnetized ion sound regime) as also was concluded in Ref. 44.

An important feature of the strong turbulence regime observed in our simulations is the difference in the ion and electron density fluctuations. While the electron density perturbations are rather benign and lower amplitude, the ion density perturbations have much larger amplitude and contain much larger short wavelength (non-quasineutral) content with $k_y \lambda_{De} \geq 1$. Such short wavelength modes do exhibit some interesting ion sound-like characteristics, such as the nonlinear ω_{pi} harmonics, nonlinear wave breaking, tendency for wave crests to become shock-like, and the elastic-like collision of wave crests without interference. It is important to note that the intense fluctuations in the short wavelength part of the spectrum are less effective in supporting the anomalous electron current as well as for the electron heating. We note that rich short wavelength features are observed in ion density but not in electron density, which is much more smooth and coherent. This difference in the electron and ion response can be important for interpretation of the fluctuation diagnostics data.⁴⁵ The coherent nature of the electron density wave and coherent electron-wave interaction is crucial for the electron heating mechanism⁴⁴ that also precludes the application of the quasilinear theory.⁴⁶

An interesting feature in our simulation is the strong $n_y = 1$ component of the axial anomalous current, resulting in the alternating jets caused by the MTSI activity. The Modified Two-Stream Instability (MTSI) that was naturally absent from the 1D simulations is shown to amplify the inverse cascade tendency due to the long wavelength nature of the MTSI even in the linear regime. Development of the Modified Two-Stream Instability results in rapid parallel heating due to the finite electric field along the magnetic field. Intense parallel heating results in increased losses of high energy electrons into the sheath that serves as an additional saturation mechanism. Similar heating was also observed in Ref. 19. Their simulations also suggest that effects of secondary emission could significantly increase anomalous transport, but unfortunately restrict wave vector space for heating studies. They too observe modulations in the sheath, however. In our simulations, the anomalous electron transport sets at the level similar to that in our 1D simulations,²¹ perhaps due to the absence of secondary emission.

SUPPLEMENTARY MATERIAL

See [supplementary material](#) for movies of three quantities discussed in the paper: δn_i , δn_e , and T_e over the simulation box $L_r \times l_\theta$. They illustrate the dynamics observed and reported in this paper.

ACKNOWLEDGMENTS

This work was supported in part by NSERC Canada and the Air Force Office of Scientific Research under Award Nos. FA9550-18-1-0132 and FA9550-15-1-0226. Computational resources from ComputeCanada/WestGrid were used in this work.

APPENDIX A: ON LINEAR GROWTH RATE ANALYSIS AND OTHER TECHNIQUES

It is possible to determine all the linear growth rates directly from the spectrogram obtained from the simulation, when the simulations are run with good resolution both in space and particle number to have the modes well resolved with $k_{\Delta x} \lambda_{De}/4 \geq 1$ and deep in the linear regime. Growth rates from the non-linear 2D simulation are obtained by discrete Fourier decomposition in space, and expressing the Fourier coefficients as a function of time

$$\phi(z, y; t) = \frac{1}{N} \sum_{k=-N/2}^{N/2} c_k(z, t) \exp iky, \quad (\text{A1})$$

and with the usual expression for a traveling wave, we have $c_k(z, t) = \tilde{c}_k(z) \exp i \int \omega(t) dt$, where $\omega(t)$ is complex-valued. Particularly in the linear regime where $\omega = \omega_r + i\gamma$ we may obtain the growth rate and frequency as a linear fit to the main branch of the complex logarithm of the Fourier coefficients as $i\omega(t) = d/dt \log(c_k)$.

Waves are largely observed to propagate in the periodic y -direction, so fluctuations of quantities are measured against the y -mean for convenience. Namely, the subtracted ‘‘average,’’ here for electron density, is defined by $\langle n_e \rangle_y(z, t) = \frac{1}{l_\theta} \int_0^{l_\theta} n_e(y, z, t) dy$. Using this density as the reference also suppresses the large drop that is observed in the sheath.

In Fig. 8, we illustrate this process as performed on our simulation data. The MTSI mode (first peak) is well represented though, showing the importance of good statistics. To emphasize this point, we ran a case for $T_e \approx 0$ with the 2D

TABLE II. Numerical parameters for the 2D ECDCI simulations.

Symbol	Value
T_e	10 eV
$n_{e,i}$	10^{17}
$\lambda_{De}/\{\Delta y, \Delta z\}$	$4/\sqrt{2}$
N_p/N_g	800
$l_\theta/\Delta y$	512
$L_r/\Delta z$	2048
B_0	0.02 T
E_0	20 kV/m
N_{cyc}	3

code, and a case with the 1D code with very good statistics (Fig. 3) that gives a very good value for both the growth rate and frequency of the Buneman instability [Eq. (18)]. In the 1D simulation, we had $N_p/N_g = 40\,000$ particles per cell and $N_g = 3400$ cells, allowing for resolution that is difficult to get with 2D simulations due to computational limitations. It would be possible to extend the growth region for ECDI by decreasing the noise level by increasing the particle number, but the simulations up to the nonlinear stage would become unfeasible. The 2D simulations use electron sub-cycling with a modest ratio of $N_{cyc} = 3$ electron steps per ion step.

- ¹I. G. Mikellides, B. Jorns, I. Katz, and A. Lopez Ortega, in 52nd AIAA/SAE/ASEE Joint Propulsion Conference, AIAA Propulsion and Energy Forum (American Institute of Aeronautics and Astronautics, 2016).
- ²T. Laffleur, S. D. Baalrud, and P. Chabert, *Phys. Plasmas* **23**, 053503 (2016).
- ³J. P. Boeuf, *J. Appl. Phys.* **121**, 011101 (2017).
- ⁴D. W. Forslund, R. L. Morse, and C. W. Nielson, *Phys. Rev. Lett.* **25**, 1266 (1970).
- ⁵S. P. Gary and J. J. Sanderson, *J. Plasma Phys.* **4**, 739 (1970).
- ⁶M. Lampe, W. Manheimer, J. B. McBride, J. H. Orens, R. Shanny, and R. N. Sudan, *Phys. Rev. Lett.* **26**, 1221 (1971).
- ⁷R. L. Stenzel and B. H. Ripin, *Phys. Rev. Lett.* **31**, 1545 (1973).
- ⁸S. Matsukiyo and M. Scholer, *J. Geophys. Res.: Space Phys.* **111**, A06104, <https://doi.org/10.1029/2005JA011409> (2006).
- ⁹L. Muschietti and B. Lembège, *J. Geophys. Res.: Space Phys.* **118**, 2267, <https://doi.org/10.1002/jgra.50224> (2013).
- ¹⁰L. Muschietti and B. Lembège, *Adv. Space Res.* **37**, 483 (2006).
- ¹¹A. Ducrocq, J. C. Adam, A. Heron, and G. Laval, *Phys. Plasmas* **13**, 102111 (2006).
- ¹²J.-P. Boeuf, *Front. Phys.* **2**, 74 (2014).
- ¹³T. Laffleur, S. D. Baalrud, and P. Chabert, *Phys. Plasmas* **23**, 053502 (2016).
- ¹⁴J. C. Adam, A. Heron, and G. Laval, *Phys. Plasmas* **11**, 295 (2004).
- ¹⁵J. Boeuf, in *The 35th International Electric Propulsion Conference*, Atlanta, GA, USA, 8–12 October 2017 (IEPC, 2017).
- ¹⁶P. Coche and L. Garrigues, *Phys. Plasmas* **21**, 023503 (2014).
- ¹⁷V. Croes, T. Laffleur, Z. Bonaventura, A. Bourdon, and P. Chabert, *Plasma Sources Sci. Technol.* **26**, 034001 (2017).
- ¹⁸K. Hara and S. Cho, in *The 35th International Electric Propulsion Conference*, Atlanta, GA, USA, 8–12 October 2017 (IEPC, 2017).
- ¹⁹A. Héron and J. C. Adam, *Phys. Plasmas* **20**, 082313 (2013).
- ²⁰J. Cavalier, N. Lemoine, G. Bonhomme, S. Tsikata, C. Honore, and D. Gresillon, *Phys. Plasmas* **20**, 082107 (2013).
- ²¹S. Janhunen, A. Smolyakov, O. Chapurin, D. Sydorenko, I. Kaganovich, and Y. Raitses, *Phys. Plasmas* **25**, 011608 (2018).
- ²²J. B. McBride, E. Ott, J. P. Boris, and J. H. Orens, *Phys. Fluids* **15**, 2367 (1972).
- ²³S. P. Gary, *J. Plasma Phys.* **4**, 753 (1970).
- ²⁴O. Buneman, *J. Nucl. Energy, Part C Plasma Phys.* **4**, 111 (1962).
- ²⁵C. N. Lashmore-Davies, *J. Phys., Part A* **3**, L40 (1970).
- ²⁶V. I. Arefev, *Sov. Phys. Tech. Phys.-USSR* **14**, 1487 (1970).
- ²⁷P. S. Foundation, *Python Programming Language* (P. S. Foundation, 2018).
- ²⁸T. E. Oliphant, *Comput. Sci. Eng.* **9**, 10 (2007).
- ²⁹V. I. Arefev, A. V. Gordeev, and L. I. Rudakov, *Nucl. Fusion* **S**, 143 (1969).
- ³⁰C. N. Lashmore-Davies and T. J. Martin, *Nucl. Fusion* **13**, 193 (1973).
- ³¹K. N. Stepanov, *Sov. Phys. Tech. Phys.* **9**, 1653 (1965).
- ³²F. F. Chen, *J. Nucl. Energy, Part C* **7**, 399 (1965).
- ³³F. F. Chen, *Phys. Fluids* **8**, 752 (1965).
- ³⁴F. F. Chen, *Phys. Fluids* **22**, 2346 (1979).
- ³⁵P. J. Barrett, R. J. Taylor, J. M. Sellen, B. D. Fried, and C. F. Kennel, *Phys. Rev. Lett.* **28**, 337 (1972).
- ³⁶D. Y. Sydorenko, “Particle-in-cell simulations of electron dynamics in low pressure discharges with magnetic fields,” Ph.D. thesis (University of Saskatchewan, 2006).
- ³⁷F. Taccogna, private communication (2017).
- ³⁸R. Davidson, *Methods in Nonlinear Plasma Theory* (Academic Press, New York, 2012).
- ³⁹M. Remoissenet, *Waves Called Solitons: Concepts and Experiments* (Springer, Berlin/Heidelberg, 2013).
- ⁴⁰A. I. Smolyakov, W. Frias, I. D. Kaganovich, and Y. Raitses, *Phys. Rev. Lett.* **111**, 115002 (2013).
- ⁴¹M. Lampe, J. B. McBride, W. Manheimer, R. N. Sudan, R. Shanny, J. H. Orens, and K. Papadopo, *Phys. Fluids* **15**, 662 (1972).
- ⁴²D. Forslund, C. Nielson, R. Morse, and J. Fu, *Phys. Fluids* **15**, 1303 (1972).
- ⁴³M. Lampe, W. Manheimer, J. B. McBride, and J. H. Orens, *Phys. Fluids* **15**, 2356 (1972).
- ⁴⁴D. W. Forslund, R. L. Morse, and C. W. Nielson, *Phys. Fluids* **15**, 2363 (1972).
- ⁴⁵S. Tsikata and T. Minea, *Phys. Rev. Lett.* **114**, 185001 (2015).
- ⁴⁶V. S. Mikhailenko, K. N. Stepanov, and E. E. Scime, *Phys. Plasmas* **10**, 2247 (2003).